CORRECTION

Open Access



Correction: PSF functions as a repressor of hypoxia-induced angiogenesis by promoting mitochondrial function

Lijie Dong^{1,2,3*†}, Wenbo Li^{1,2,3}, Tingting Lin^{1,2,3}, Boshi Liu^{1,2,3}, Yaru Hong^{1,2,3}, Xiaomin Zhang^{1,2,3} and Xiaorong Li^{1,2,3*}

Correction: Cell Commun Signal 19, 14 (2021) https://doi.org/10.1186/s12964-020-00684-w

Following publication of the original article [1], the authors reported an error in Fig. 5b and c.

Due to an error in the manuscript preparation and proofreading stage, there are mistakes in Fig. 5b and c, including image misuse (Fig. 5b) and incorrect labeling (Fig. 5b and c). The figure presented in this correction article has been corrected.

Published online: 24 July 2023

[†]Lijie Dong: the frst author of this manuscript.

The original article can be found online at https://doi.org/10.1186/s12964-020-00684-w

*Correspondence: Lijie Dong aitaomubang@126.com Xiaorong Li xiaorli@163.com ¹ Tianjin Key Laboratory of Retinal Functions and Diseases, Tianjin, People's Republic of China ² Tianjin International Joint Research and Development Centre of Ophthalmology and Vision Science, Tianjin, People's Republic of China ³ Eye Institute and School of Optometry, Tianjin Medical University Eye Hospital, 251 Fukang Road, Nankai, Tianjin 300384, People's Republic of China



© The Author(s) 2023. Open Access This article is licensed under a Creative Commons Attribution 4.0 International License, which permits use, sharing, adaptation, distribution and reproduction in any medium or format, as long as you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons licence, and indicate if changes were made. The images or other third party material in this article are included in the article's Creative Commons licence, unless indicated otherwise in a credit line to the material. If material is not included in the article's Creative Commons licence and your intended use is not permitted by statutory regulation or exceeds the permitted use, you will need to obtain permission directly from the copyright holder. To view a copy of this licence, visit http://creativecommons.org/licenses/by/4.0/. The Creative Commons Public Domain Dedication waiver (http://creativeco mmons.org/publicdomain/zero/1.0/) applies to the data made available in this article, unless otherwise stated in a credit line to the data

Reference

1. Dong L, Li W, Lin T, et al. PSF functions as a repressor of hypoxia-induced angiogenesis by promoting mitochondrial function. Cell Commun Signal. 2021;19:14. https://doi.org/10.1186/s12964-020-00684-w.



Blot:anti-HIF-1α

Fig. 5 b-c PSF-HIF-1α complex formation under hypoxia induction. Ectopically expressed PSF interacted with HIF-1α were checked using anti-HA antibody in HRMECs as indicated (**a**) and anti-His antibody sedimentation (**b**). **c** Anti-HIF-1αor anti-PSF antibody-based immunoprecipitation with total cell lysates of HRMECs under hypoxia exposure was performed, followed by immunoblotting with the corresponding antibodies as indicated