

Meeting abstract

Calcium signals in lymphocyte activation and disease

S Feske

Address: NYU School of Medicine, Department of Pathology, New York, USA
from 12th Joint Meeting of the Signal Transduction Society (STS). Signal Transduction: Receptors, Mediators and Genes
Weimar, Germany. 29–31 October 2008

Published: 26 February 2009

Cell Communication and Signaling 2009, **7**(Suppl 1):A76 doi:10.1186/1478-811X-7-S1-A76

This abstract is available from: <http://www.biosignaling.com/content/7/S1/A76>

© 2009 Feske; licensee BioMed Central Ltd.

Calcium ions function as universal second messengers in virtually all eukaryotic cells including cells of the immune system where they are crucial for the function of T and B cells, mast cells and dendritic cells. The predominant mechanism regulating intracellular Ca^{2+} levels in cells of the adaptive immune system is store-operated Ca^{2+} influx through so-called Ca^{2+} -release activated Ca^{2+} (CRAC) channels. We identified ORAI1 (also named CRACM1) as a pore subunit of the CRAC channel essential for the function of T cells and mast cells. ORAI1/CRAC channels are activated when intracellular Ca^{2+} stores are depleted. The resulting decrease in the ER Ca^{2+} concentration is sensed by stromal interaction molecule 1 (STIM1) that is required for activation of ORAI1/CRAC channels. We showed that murine T cells lacking STIM1 exhibit severely impaired store-operated Ca^{2+} influx. T cells from mice lacking STIM1 or its paralogue STIM2 both showed significantly reduced cytokine production *in vitro* and a defect in regulatory T cell development as well as lympho- and myeloproliferation *in vivo*. Mutation of ORAI1 in humans is associated with severe combined immunodeficiency (SCID), increased susceptibility to infections and a failure to thrive. A similar defect is found in mice transgenic for the equivalent R93W mutation in murine ORAI1, which all but abrogates CRAC channel function and T cell activation. Taken together STIM1, STIM2 and ORAI1 are essential regulators of store-operated Ca^{2+} entry in cells of the immune system and other tissues.