

SHORT REPORT

Open Access

Src mediates cytokine-stimulated gene expression in airway myocytes through ERK MAPK

Cherie A Singer^{1*}, Beata Lontay¹, Helmut Unruh², Andrew J Halayko³ and William T Gerthoffer^{1,4}

Abstract

The p38 and extracellular signal-regulated kinases (ERK) mitogen-activated protein kinases (MAPK) participate in cytokine-stimulated inflammatory gene expression in airway smooth muscle cells. The following study was undertaken to determine whether Src tyrosine kinases are signaling intermediaries upstream of cytokine-stimulated MAPK activation and gene expression. Treating human airway myocytes with interleukin (IL)-1 β , tumor necrosis factor (TNF) α and interferon (IFN) γ caused a rapid 1.8-fold increase in Src family tyrosine kinase activity within 1 minute that remained 2.3 to 2.7 fold above basal conditions for 15 minutes. This activity was blocked by addition of 30 μ M PP1, a pyrimidine inhibitor specific for Src family tyrosine kinases, in immune-complex assays to confirm that this stimulus activates Src tyrosine kinase. Addition of PP1 also blocked cytokine-stimulated expression of IL-1 β , IL-6 and IL-8, while decreasing phosphorylation of ERK, but not p38 MAPK. Since this inflammatory stimulus may activate additional inflammatory signaling pathways downstream of Src, we tested the effects of PP1 on phosphorylation of signal transducers and activators of transcription (STAT). PP1 had no effect on cytokine-stimulated STAT 1 or STAT 3 phosphorylation. These results demonstrate that Src tyrosine kinases participate in the regulation of IL-1 β , IL-6 and IL-8 expression and that these effects of Src are mediated through activation of ERK MAPK and not p38 MAPK or STAT1/STAT3 phosphorylation.

Findings

Our laboratory has examined signaling pathways regulating secretion of inflammatory mediators by human airway smooth muscle cells. The synthesis and secretion of Th1/Th2 cytokines, along with CC and C-X-C chemokines, chemotactic proteins, peptide growth factors and their receptors can be induced in these myocytes by exposure to, among others, interleukin (IL)-1 β , tumor necrosis factor (TNF) α , interferon (IFN) γ , or transforming growth factor β [1,2] and contributes to inflammatory airway disease. In previous studies, we used a complementary DNA expression array to analyze expression of inflammatory mediators following treatment with a pro-inflammatory stimulus consisting of IL-1 β , TNF α and IFN γ and established that this stimulus induces expression of multiple inflammatory mediators including IL-1 β , IL-6, and IL-8 [3]. Pharmacological inhibitors of mitogen-activated protein kinase (MAPK) activation were used to further demonstrate that both

p38 and ERK MAPK are upstream mediators of IL-6 and IL-8 expression, while ERK MAPK alone was involved in mediating IL-1 β expression.

Src tyrosine kinases are one of the signaling intermediaries linking multiple types of receptors to MAPK activation in smooth muscle. In colonic myocytes, the G-protein coupled M₂ muscarinic receptor is coupled to ERK MAPK through Src activation [4] and expression of IL-1 β , IL-6, IL-8, and cyclooxygenase (COX-2) mRNA is reduced by inhibition of Src with the pyrimidine inhibitor PP1 [5]. In vascular smooth muscle cells, angiotensin II stimulates Src-dependent p38 MAPK activation [6] and CD40 ligation initiates Src activation of p38 and ERK MAPK, resulting in the induction of IL-8 and monocyte chemotactic protein-1 [7]. In airway myocytes, the PDGF receptor signals to ERK MAPK through Src activation via a pertussis-toxin sensitive mechanism that suggests the involvement of G_i-protein subunits [8] and sphingosine-1-phosphate stimulation of a G_i-coupled receptor also stimulates ERK MAPK through Src [9]. In addition to acting as an upstream mediator of MAPK, Src may also activate other signaling pathways that could affect inflammatory gene expression,

* Correspondence: csinger@medicine.nevada.edu

¹University of Nevada School of Medicine, Department of Pharmacology
Reno, NV 89557, USA

Full list of author information is available at the end of the article


such as the Janus kinase-signal transducers and activators of transcription (JAK-STAT). This is supported by evidence that inhibition of Src in PDGF-stimulated airway myocytes with PP2 blocks both ERK MAPK and JAK2 activation, resulting in decreased phosphorylation of STAT1 and STAT3 [10] that requires an interaction with the small GTPase Rac1 [11].


The following study was undertaken to determine whether activation of Src is an event upstream of MAPK signaling contributing to cytokine-stimulated gene expression and all methodological details are described in Additional File 1. *In vitro* kinase assays initially evaluated the activity of Src in human airway myocytes isolated according to previously published protocols using p34^{cdc2} (6-20) as the substrate. This substrate is specific for Src tyrosine kinase family members including Src, Lck, Lyn and Fyn but not non-Src tyrosine kinases such as Abl [12]. Stimulation of cultures with 10 ng/ml IL-1 β , TNF α and IFN γ caused a rapid 1.8-fold increase in Src family tyrosine kinase activity within 1 minute that remained 2.3 to 2.7-fold above basal conditions for 15 minutes (Figure 1A) and addition of the inhibitor PP1 prevented cytokine stimulation of kinase activity. This inhibitor, and the similar compound PP2, have been widely used to pharmacologically inhibit multiple Src family members [13-15], and PP1 can also inhibit Abl, which was presumably not measured in our assay with the p34^{cdc2} (6-20) substrate. To confirm the activity measured was attributed to Src, we performed additional immune-complex kinase assays using the p34^{cdc2} (6-20) substrate following immunoprecipitation with a Src-specific antibody, as

described [4]. In these assays (Figure 1B), a 2-fold increase in activity of Src immunoprecipitates within 5 minutes of stimulation was abolished by addition of 30 μ M PP1, indicative of Src-specific activity in airway myocytes.

To determine whether Src activation contributes to inflammatory gene expression, we chose to evaluate the expression of genes previously shown to be regulated by p38 and ERK MAPK, namely IL-1 β , IL-6 and IL-8 [3]. Gene expression was measured using a relative semi-quantitative RT-PCR approach in which the target gene is amplified within the linear range in a multi-plex reaction containing 18S rRNA as an endogenous standard. We further verified the results of these experiments by quantifying synthesis of the target protein by Western blots or ELISA. As shown previously [3], expression of IL-1 β , IL-6 and IL-8 is induced upon stimulation with 10 ng/ml IL-1 β , TNF α and IFN α for 20 hours (Figure 2). Treatment with 30 μ M PP1 reduces cytokine-stimulated IL-1 β mRNA expression by 53% (Figure 2A), which correlates by a 64% decrease in IL-1 β synthesis, as measured by immunoblotting for the intracellular 33 kDa form of IL-1 β (Figure 2D). This was necessary since high levels of IL-1 β present in the media from the cytokine-stimulus would hamper results from ELISA. Treatment with PP1 also inhibited IL-6 mRNA expression by 43% (Figure 2B) and IL-8 mRNA expression by 56% (Figure 2C), which again corresponded to a 80% decrease in IL-6 secretion from 160.8 ng/ml to 32.9 ng/ml (Figure 2E) and a 82% decrease in IL-8 secretion from 155.6 ng/ml to 27.2 ng/ml (Figure 2F). Thus, the same concentration of PP1 that effectively blocks Src kinase activity results in inhibition, but not complete blockade, of inflammatory gene expression. This suggests that while Src activity is involved in expression of the genes examined, additional Src-independent signaling pathways likely participate.

To determine whether Src is an upstream mediator of MAPK activation in airway myocytes, we analyzed MAPK activation by phospho-immunoblot analysis in cultures treated with PP1. Treatment with 10 ng/ml IL-1 β , TNF α and IFN γ for 15 minutes resulted in a 10.2 fold increase in ERK MAPK phosphorylation that was inhibited 50% by addition of PP1 (Figure 3A). The same stimulus resulted in a 2.5 fold increase in p38 MAPK phosphorylation but addition of PP1 had no effect (Figure 3B). Since previous reports demonstrated that PDGF stimulation of airway myocytes results in Src-dependent phosphorylation of STAT1/STAT3 via JAK2 activation [10], we determined whether the inflammatory stimulus used here may activate JAK-STAT signaling pathways to mediate downstream gene expression (Figure 3C). The combination of 10 ng/ml IL-1 β , TNF α and IFN γ increased STAT1 phosphorylation 2.5 fold and STAT3 phosphorylation 3.7 fold above basal levels






within 10 minutes. However, addition of 30 μ M PP1 did not affect STAT phosphorylation, indicating that this with this inflammatory stimulus, Src does not mediate JAK-STAT signaling.

Clearly, the functions of Src tyrosine kinases in smooth muscle cells are varied and likely involve interactions with multiple signaling pathways. In the airway, Src mediates serotonin-evoked peak Ca²⁺ responses by affecting phosphoinositide levels to alter cell contraction [15]. Src also mediates PDGF and thrombin-induced proliferative responses, possibly through stimulation of cyclin D1 expression [10,13]. While previous studies have established that both ERK and p38 MAPK are involved in regulating expression of many inflammatory genes in airway myocytes [3,16,17], this work demonstrates that Src tyrosine kinases also play a role in regulating inflammatory gene expression by signaling upstream of ERK but not p38 MAPK. This is supported

by studies of inflammatory gene expression in other cell types. In pulmonary epithelial cells, both ERK and p38 MAPK contribute to silica-induced IL-8 release but only ERK MAPK activation is dependent on Src [14]. CD40 stimulation of IL-8 and MCP-1 production in vascular myocytes is also dependent on ERK and p38 MAPK activation but the inhibitor PP2 was found to inhibit only ERK MAPK [7]. In this same study, PP2 also decreased activation of I κ B kinase, indicating that Src stimulates NF- κ B signaling to affect chemokine production and suggesting that NF- κ B activation could link cytokine stimulation to MAPK activation. We have previously shown that IL-1 β and TNF α but not IFN γ activates NF- κ B in airway myocytes and inhibition of NF- κ B activity reduces expression of IL-1 β , IL-6, IL-8 and COX-2 in a manner independent of p38 MAPK [5,16].

The link between inflammatory signaling and Src-dependent MAPK activation in our studies remains to



be determined. One possibility may be the TNF receptor-associated proteins (TRAFs). Binding of IL-1 β to the IL-1 receptor activates IL-1 receptor associated kinases that recruit TRAFs. TNF receptors also recruit TRAFs through the TNF receptor DEATH domain (TRADD). In fibroblasts, Src interacting with TRAF2 links TNF α stimulation to ERK MAPK [18] and TRAF proteins also activate NF- κ B through recruitment of I κ B kinase (see review by [19]). The contribution of IFN γ signaling to Src-dependent ERK MAPK activation is less clear. Interactions between Src tyrosine kinases and receptor-associated JAKs are often required for complete STAT activation [20] but downstream activation of ERK MAPK has not been demonstrated. It has been proposed that IFN γ can directly activate ERK MAPK pathways through MEKK1 [21] but the signaling intermediates, if any, are unknown. Another possibility could involve heterotrimeric G $_i$ subunits implicated in Src-dependent activation of ERK MAPK in airway myocytes stimulated with PDGF [8]. The contribution of these proteins have not been widely examined in the context of IL-1 β TNF α or IFN γ signaling, which are more likely to utilize small GTPases such as Rac1 [11] and RhoA [22]. Thus, while these results demonstrate that Src participate in the regulation of inflammatory gene expression through activation of ERK MAPK, further studies will explore the signaling pathways linking Src-dependent cytokine stimulation to MAPK activation.

Additional material

Additional file 1: The file provided detailed methods described in the communication [23].

Acknowledgements and Funding

The authors would like to thank Ms. Shanti Rawat and Ms. Michelle Deetken for their excellent technical assistance. This work was supported by a Minority Investigator Research Supplement from the NIH/NHLBI to CAS and by the NIH/NHLBI HL48183-10 to W.T.G.

Author details

¹University of Nevada School of Medicine, Department of Pharmacology Reno, NV 89557, USA. ²University of Manitoba, Section of Thoracic Surgery, Winnipeg, MB, Canada R3A 1R9. ³University of Manitoba, Department of Physiology and Section of Respiratory Diseases, Winnipeg, MB, Canada R3A 1R8. ⁴University of South Alabama, Department of Biochemistry & Molecular Biology, Mobile, AL 36688, USA.

Authors' contributions

CAS and BL performed all the experiments. HU and AJH provided and characterized human cells used in the experiments. CAS wrote the manuscript and AJH and W.T.G. provided comments for revision. All authors have read and approved the final manuscript.

Competing interests

The authors declare that they have no competing interests.

Received: 5 January 2011 Accepted: 20 May 2011

Published: 20 May 2011

References

- Singer CA, Salinithone S, Baker KJ, Gerthoffer WT: **Synthesis of immune modulators by smooth muscles.** *Bioessays* 2004, **26**:646-655.
- Knox AJ, Pang L, Johnson S, Hamad A: **Airway smooth muscle function in asthma.** *Clin Exp Allergy* 2000, **30**:606-614.

3. Hedges JC, Singer CA, Gerthoffer WT: **Mitogen-activated protein kinases regulate cytokine gene expression in human airway myocytes.** *Am J Respir Cell Mol Biol* 2000, **23**:86-94.
4. Singer CA, Vang S, Gerthoffer WT: **Coupling of M(2) muscarinic receptors to Src activation in cultured canine colonic smooth muscle cells.** *Am J Physiol Gastrointest Liver Physiol* 2002, **282**:G61-G68.
5. Salinthon S, Singer CA, Gerthoffer WT: **Inflammatory gene expression by human colonic smooth muscle cells.** *Am J Physiol Gastrointest Liver Physiol* 2004.
6. Touyz RM, He G, El MM, Diep Q, Mardigyan V, Schiffrin EL: **Differential activation of extracellular signal-regulated protein kinase 1/2 and p38 mitogen activated-protein kinase by AT1 receptors in vascular smooth muscle cells from Wistar-Kyoto rats and spontaneously hypertensive rats.** *J Hypertens* 2001, **19**:553-559.
7. Mukundan L, Milhorn DM, Matta B, Suttles J: **CD40-mediated activation of vascular smooth muscle cell chemokine production through a Src-initiated, MAPK-dependent pathway.** *Cell Signal* 2004, **16**:375-384.
8. Conway AM, Rakhit S, Pyne S, Pyne NJ: **Platelet-derived-growth-factor stimulation of the p42/p44 mitogen-activated protein kinase pathway in airway smooth muscle: role of pertussis-toxin-sensitive G-proteins, c-Src tyrosine kinases and phosphoinositide 3-kinase.** *Biochem J* 1999, **337**:171-177.
9. Rakhit S, Conway AM, Tate R, Bower T, Pyne NJ, Pyne S: **Sphingosine 1-phosphate stimulation of the p42/p44 mitogen-activated protein kinase pathway in airway smooth muscle. Role of endothelial differentiation gene 1, c-Src tyrosine kinase and phosphoinositide 3-kinase.** *Biochem J* 1999, **338**(Pt 3):643-649.
10. Simon AR, Takahashi S, Severgnini M, Fanburg BL, Cochran BH: **Role of the JAK-STAT pathway in PDGF-stimulated proliferation of human airway smooth muscle cells.** *Am J Physiol Lung Cell Mol Physiol* 2002, **282**:L1296-L1304.
11. Simeone-Penney MC, Severgnini M, Rozo L, Takahashi S, Cochran BH, Simon AR: **PDGF-induced human airway smooth muscle cell proliferation requires STAT3 and the small GTPase Rac1.** *Am J Physiol Lung Cell Mol Physiol* 2008, **294**:L698-704.
12. Cheng HC, Nishio H, Hatase O, Ralph S, Wang JH: **A synthetic peptide derived from p34cdc2 is a specific and efficient substrate of src-family tyrosine kinases.** *J Biol Chem* 1992, **267**:9248-9256.
13. Tsang F, Choo HH, Dawe GS, Wong WS: **Inhibitors of the tyrosine kinase signaling cascade attenuated thrombin-induced guinea pig airway smooth muscle cell proliferation.** *Biochem Biophys Res Commun* 2002, **293**:72-78.
14. Orevik J, Lag M, Schwarze P, Refsnes M: **p38 and Src-ERK1/2 pathways regulate crystalline silica-induced chemokine release in pulmonary epithelial cells.** *Toxicol Sci* 2004, **81**:480-490.
15. Tolloczko B, Turkewitsch P, Choudry S, Bisotto S, Fixman ED, Martin JG: **Src modulates serotonin-induced calcium signaling by regulating phosphatidylinositol 4,5-bisphosphate.** *Am J Physiol Lung Cell Mol Physiol* 2002, **282**:L1305-L1313.
16. Singer CA, Baker KJ, McCaffrey A, AuCoin DP, Dechert MA, Gerthoffer WT: **p38 MAPK and NF-κB mediate COX-2 expression in human airway myocytes.** *Am J Physiol Lung Cell Mol Physiol* 2003, **285**:L1087-L1098.
17. Laporte JD, Moore PE, Abraham JH, Maksym GN, Fabry B, Panettieri RA Jr, Shore SA: **Role of ERK MAP kinases in responses of cultured human airway smooth muscle cells to IL-1β.** *Am J Physiol* 1999, **277**:L943-L951.
18. van Vliet C, Bukczynska PE, Puryer MA, Sadek CM, Shields BJ, Tremblay ML, Tiganis T: **Selective regulation of tumor necrosis factor-induced Erk signaling by Src family kinases and the T cell protein tyrosine phosphatase.** *Nat Immunol* 2005, **6**:253-260.
19. Lee NK, Lee SY: **Modulation of life and death by the tumor necrosis factor receptor-associated factors (TRAFs).** *J Biochem Mol Biol* 2002, **35**:61-66.
20. Garcia R, Bowman TL, Niu G, Yu H, Minton S, Muro-Cacho CA, Cox CE, Falcone R, Fairclough R, Parsons S, Laudano A, Gazit A, Levitzki A, Kraker A, Jove R: **Constitutive activation of Stat3 by the Src and JAK tyrosine kinases participates in growth regulation of human breast carcinoma cells.** *Oncogene* 2001, **20**:2499-2513.
21. Roy SK, Hu J, Meng Q, Xia Y, Shapiro PS, Reddy SP, Platanias LC, Lindner DJ, Johnson PF, Pritchard C, Pagés G, Pouyssegur J, Kalvakolanu DV: **MEKK1 plays a critical role in activating the transcription factor C/EBP-beta-dependent gene expression in response to IFN-γ.** *Proc Natl Acad Sci USA* 2002, **99**:7945-7950.
22. Hunter I, Nixon GF: **Spatial compartmentalization of tumor necrosis factor (TNF) receptor 1-dependent signaling pathways in human airway smooth muscle cells. Lipid rafts are essential for TNF-α-mediated activation of RhoA but dispensable for the activation of the NF-κB and MAPK pathways.** *J Biol Chem* 2006, **281**:34705-34715.
23. Rahman MS, Shan LY, Yang J, Unruh H, Yang X, Halayko AJ, Gounni AS: **IL-17R activation of human airway smooth muscle cells induces CXCL-8 production via transcriptional dependent mechanism.** *Clin Immunol* 2005, **115**:268-276.

doi:10.1186/1478-811X-9-14

Cite this article as: Singer et al.: Src mediates cytokine-stimulated gene expression in airway myocytes through ERK MAPK. *Cell Communication and Signaling* 2011 **9**:14.

Submit your next manuscript to BioMed Central and take full advantage of:

- Convenient online submission
- Thorough peer review
- No space constraints or color figure charges
- Immediate publication on acceptance
- Inclusion in PubMed, CAS, Scopus and Google Scholar
- Research which is freely available for redistribution

Submit your manuscript at
www.biomedcentral.com/submit

